

A Geographically and Seasonally Resolved Ammonia Emission Inventory for Dairy Farms

Robert W. Pinder, Natalie J. Anderson, Ross Strader, Cliff I.
Davidson, Peter J. Adams

Carnegie Mellon University

12th International Emission Inventory Conference
San Diego, CA
April 30, 2003


Ammonia

- Contributes to formation of inorganic particulate matter
- Deposition of Nitrogen
 - Forest ecosystems
 - Surface and coastal waters
- Not a criteria pollutant under Clean Air Act
- Sources:
 - 55-85% volatilization from livestock manure
 - Dairy farms are substantial fraction: 20-60%

Seasonal and Geographic Variation

- Ammonium nitrate aerosol is a significant fraction of $\text{PM}_{2.5}$
- Sensitivity of $[\text{PM}]$ to $[\text{NH}_3]$
 - Dependent on
 - Temperature
 - Relative Humidity
 - $[\text{HNO}_3^-]$
 - Ranges from 0 to 4.7

Uncertainty and Variability in Emissions




- Variability
 - Farming practices
 - Climate
- Uncertainty
 - Incomplete understanding
 - Measurement error

Source: Plochl, Matthias. Neural network approach for modelling ammonia emission after manure application on the field, *Atmospheric Environment*. 35 (2001) 5833-5841


Approach

- Partially mechanistic model of ammonia emissions from a single dairy farm
 - Combines experimental results and physical principles to explain variation in emission factors
 - Focus on temporal and geographic variation
- Combine farm model with farming practices to generate national inventory
- Tool for evaluating change in emissions from regulation



Farm Emissions Model

Nitrogen Flows



Inputs:

From Housing:
Volume
TAN mass
Urea mass

Climate:
Temperature
Wind Speed
Precipitation
pH
Mass transfer
resistance

Storage Sub-Model

Variables:

- Volume

$$dV/dt = \text{precipitation} - \text{evaporation}$$

- Mass of urea nitrogen

$$dU/dt = - \text{urea hydrolysis rate law}$$

- Mass of Total Ammonia Nitrogen (TAN)

$$dM/dt = \text{urea loading} - \text{emissions}$$

$$\text{Emissions} = \text{Area}[\text{TAN}]H^*r^{-1}$$

H^* : Effective Henry's Law Constant

r : mass transfer resistance

Outputs: NH_3 Volatilized
To Application: Volume
Mass TAN

Reported Inputs:
Values as reported


T: 8 12 15

U_{10} : 1.3 m/sec

Unreported Inputs:
Ranges based on similar research


pH: 7.9 8.3 8.5

Prior Distribution for Parameter:



Farm Emissions Model:


Calculate likelihood for each iteration



Likelihood = $\Pr(\text{model} \mid \text{data})$


Update Posterior Distribution

Posterior Distribution for Parameter:



Verification of Tuned Parameters


Storage Surface Resistance




Tuned Resistance Parameters

Storage Surface Resistance

No crust




Wheat Straw Cover




National Practices Model

- Survey data from USDA
- Data are sparse
- Stepwise logistic regression to predict 13 different farming practices
- Predictors:
 - Temperature, Rainfall
 - Historical cow populations
 - Milk yield
 - Other practices on the farm


Verification of Farming Practices




National Inventory




Seasonal Variability in National Emissions






January




July

kg NH₃ cow⁻¹ month⁻¹



Ratio of July to January Emission Factors



Evaluate Regulatory Options

- Substitute traditional farming practices with control strategies
- Apply proportionally with National Practices Model

Control Strategy	Percent Reduced	Range
Wheat Straw Cover Only	4.5%	3.3-4.6%
Injection Only	27%	25%-31%
Both Straw & Injection	41%	37%-43%

Conclusions

- Seasonally and geographically varied emission inventory
 - Factor of 4 seasonal change
 - As high as factor of 7 geographical change
- Reducing emissions
 - In the winter, reduce housing emissions
 - In the spring and fall, reduce application emissions
 - Need to address entire farm
- Use air quality model to validate

Questions and Comments